arXiv:astro-ph/0111347v1 17 Nov 2001

# X-ray emission from a merger remnant, NGC 7252,the "Atoms-for-Peace" galaxy

Hisamitsu Awaki

*Department of Physics, Faculty of Science, Ehime University*

Hironori Matsumoto[1]

*Department of Earth and Space Science, Osaka University,*

and

Hiroshi Tomida

*Space Utilization Research Program, National Space Development Agency of Japan,*

## ABSTRACT

We observed a nearby merger remnant NGC 7252 with the X-ray satellite *ASCA*, and detected X-ray emission with the X-ray flux of $(1.8\pm0.3)\times10^{-13}$ ergs $\mathrm{s}^{-1}$ $\mathrm{cm}^{-2}$ in the 0.5–10 keV band. This corresponds to the X-ray luminosity of $8.1\times10^{40}$ ergs $\mathrm{s}^{-1}$. The X-ray emission is well described with a two-component model: a soft component with $kT = 0.72\pm0.13$ keV and a hard component with $kT > 5.1$ keV. Although NGC 7252 is referred to as a dynamically young protoelliptical, the 0.5–4 keV luminosity of the soft component is about $2\times10^{40}$ ergs $\mathrm{s}^{-1}$, which is low for an early-type galaxy. The ratio of $L_{\mathrm{X}}/L_{\mathrm{FIR}}$ suggests that the soft component originated from the hot gas due to star formation. Its low luminosity can be explained by the gas ejection from the galaxy as galaxy winds. Our observation reveals the existence of hard X-ray emission with the 2–10 keV luminosity of $5.6\times10^{40}$ ergs $\mathrm{s}^{-1}$. This may indicate the existence of nuclear activity or intermediate-mass black hole in NGC 7252.

*Subject headings:* galaxies: evolution — galaxies: individual (NGC 7252) — X-rays: galaxies

---

[1] Center for Space Research, Massachusetts Institute of Technology, 77 Massachusetts Avenue, NE80-6045, Cambridge, MA02139-4307, USA

## 1. Introduction

NGC 7252 is a nearby galaxy (63.2 Mpc) with tidal tails in the optical band. The tidal tails are thought to be the result of a galaxy merger of two spiral galaxies (e.g. Mihos, Dubinski & Hernquist 1998). The merger phenomenon probably induced a violent starformation in the whole body (e.g. Hibbard et al. 1994, hereafter H94). We can observe traces of the violent starformation which occurred less than 1 Gyr ago; for instance, the presence of an "E+A" spectrum (Schweitzer 1982), the absence of H I gas in the inner region (Hibbard & van Gorkom 1996), and the existence of young globular clusters (e.g. Whitemore et al. 1997). In the central region, its surface profile is well represented by the $r^{1/4}$ law which is seen in early-type galaxies (e.g. Schweizer 1982, Stanford & Bushouse 1991). Numerical simulations predict that the tidal tails will disappear within a few Gyr, and then this galaxy will be recognized as an early-type galaxy (e.g. Hibbard & Mihos 1995). Therefore, this galaxy is also called "a dynamically young protoelliptical".

Early-type galaxies sometimes have powerful X-ray emission which originates from a hot gaseous halo surrounding the galaxies. Their X-ray luminosities range from $10^{39}$ to $10^{42}$ ergs $\mathrm{s}^{-1}$ (e.g. Forman, Jones, & Tucker 1985; Canizares, Fabbiano, & Trinchieri 1987). The total amount of the hot gaseous halo and the metal abundance are estimated to be $10^8$–$10^{11} M_\odot$ and sub-solar, respectively (e.g. Awaki et al. 1994; Matsushita, Ohashi, & Makishima 2000). In order to reveal the evolution of the hot gas, many observational and theoretical works have been performed. Since NGC 7252 is evolving into an early-type galaxy, this galaxy is one of the best objects for investigating the evolution of the hot gas surrounding early-type galaxies. H94 detected X-rays from NGC 7252 with the *ROSAT* PSPC and HRI, and found that the X-ray emitting region is marginally extended, which indicates the existence of hot gas in NGC 7252. However, they could not determine its accurate spectral shape. Using the *ASCA*'s wide-band spectroscopy, we reveal the nature of the hot gas in NGC 7252.

## 2. Observation and Results

NGC 7252 was observed with *ASCA* on 1998 November 16–18 (Tanaka, Inoue, & Holt 1994). Since the intense shower of Leonid meteors was expected to appear on November 17, the high voltage of the GIS was turned off in order to minimize the impact to the GIS. Therefore, the galaxy was observed with only the solid-state imaging spectrometer (SIS). The SIS consists of two detectors, called SIS0 and SIS1. The SIS data were taken in the 1 CCD FAINT mode at all bit-rates. The data reduction was performed by the *ASCA* Guest Observatory at NASA/GSFC using the following main criteria: (1) the angle from bright Earth's limb $> 20°$, (2) the elevation angle from Earth's limb $> 10°$, (3) the geometrical cut off rigidity $> 6$ GeV $\mathrm{c}^{-1}$. The detail of the selection criteria is provided from the *ASCA* Guest Observatory. The on-source exposure time after the above screening is about 43.5 ks.

We obtained an X-ray image in the 0.5–10 keV band with the data from SIS0 and SIS1. Figure 1 shows the X-ray contour map overlaid on the digitized sky survey map of NGC 7252. We accumulated SIS0 and SIS1 data within a $6'$ diameter circle centered on the X-ray source; a background spectrum is taken from a non-source region $> 8'$ in diameter. We combine energy bins to contain more than 50 counts in each bin and apply a $\chi^2$ method in the XSPEC 11.00 package for spectral fitting. The spectrum after background subtraction is shown in Figure 2.

We first attempted to fit a single-component model, a thermal model or a power law model, to the spectrum. However, the spectrum was characterized with $kT \sim$2.3 keV or $\Gamma \sim$2.3 with $\chi^2_\nu > 1.7$. Next, we introduced a two-component model, thin thermal plus thermal bremsstrahlung model. This model provided an acceptable fit to the data ( $\chi^2_\nu \sim 1.2$), with the absorption column fixed at the Galactic value of $2\times10^{20}$ $\mathrm{cm}^{-2}$ (Stark et al. 1990). Since the confidence region of the metal abundance $A_{\mathrm{RS}}$ was too large, we fixed $A_{\mathrm{RS}}$ at one solar metal abundance (Table 1). The thin thermal and the thermal bremsstrahlung models are characterized with $kT = 0.72\pm0.13$ keV and $kT > 5.1$ keV, respectively. The best-fit model is shown in Figure 2. Using this model, the X-ray flux is $(1.8\pm0.3)\times10^{-13}$ ergs $\mathrm{s}^{-1}$ $\mathrm{cm}^{-2}$ in the 0.5–10 keV band, which corresponds to the X-ray luminosity of $8.1\times10^{40}$ ergs $\mathrm{s}^{-1}$ assuming the distance of 63.2 Mpc. The X-ray flux in the *ROSAT* band (0.2–4 keV) is about $1.0\times10^{-13}$ ergs $\mathrm{s}^{-1}$ $\mathrm{cm}^{-2}$, which is consistent with that obtained by H94, although they did not determine the spectral shape of the X-rays from NGC 7252, and, hence assumed a thin thermal emission with $kT = 1$ keV and metal abundance of 1 solar.

## 3. Discussion

### 3.1. Origin of the Soft Component

The *ROSAT* observation by H94 found a soft X-ray emission which is marginally extended, which indicates the existence of the hot gas in NGC 7252. Our *ASCA* observation finds that the soft X-ray emission has the characteristic temperature of $kT = 0.72 \pm 0.13$ keV, and the observed X-ray flux in the 0.5–4 keV band is about $4 \times 10^{-14}$ ergs s$^{-1}$ cm$^{-2}$, which corresponds to the X-ray luminosity of the $2 \times 10^{40}$ ergs s$^{-1}$ after the correction of the Galactic absorption. As compared with X-ray properties of early type galaxies, the soft X-ray luminosity of NGC 7252 is low for an early-type galaxy. Our result is consistent with a claim by Fabbiano & Schweizer (1995), who pointed out that NGC 7252 belongs to group 1 in the $L_{\rm B} - L_{\rm X}$ diagram of E and S0 galaxies (see figure 7 in their paper). In group 1, the contribution of the soft component to the total X-ray emission is considered to be small (e.g. Fabbiano & Schweizer 1995, Matsumoto et al. 1997).

NGC 7252 is a post-starburst galaxy as well as a dynamically young protoelliptical (e.g. Schweizer 1982, Stanford & Bushouse 1991). Both types of galaxies sometimes have hot gas surrounding the galaxy. We took the luminosity ratio of $L_{\rm X}/L_{\rm FIR}$ to find a possible origin of the hot gas in the galaxy. Figure 3 shows the correlation plot of the X-ray luminosities in the 0.5–4 keV band and the far-infrared luminosities. The region between two solid lines in Figure 3 indicates the luminosity ratio of about $10^{-4}$ which most starburst/H II galaxies have ( David, Jones, & Forman (1992)). The open circles in Figure 3 show the data for early-type galaxies. Their X-ray data are taken from Canizares et al. (1987), and their infrared data are deduced from IRAS 60 and 100 $\mu$m fluxes (taken from Knapp et al. (1989)) using the usual "bolometric" far-infrared brightness (e.g. Devereux & Eales 1989)

$$F_{\rm FIR} = 1.24 \times 10^{-14} (2.58 F_{60 \mu {\rm m}} + F_{100 \mu {\rm m}}) \ {\rm erg\ cm^{-2}\ s^{-1}}\ , \quad (1)$$

where the 60 and 100 $\mu$m flux densities are expressed in mJy. The far-infrared light from early-type galaxies is emitted from cool interstellar matter, and thus their luminosities are smaller than those of starburst galaxies (Knapp et al. 1989; Bregman et al. 1998). Therefore, early-type galaxies lie in the upper left side in the figure. NGC 7252 is a luminous far-infrared object with $L_{\rm FIR} \sim 4 \times 10^{10} L_{\odot}$ (Dupraz et al. 1990). We find that the luminosity ratio of $L_{\rm X}/L_{\rm FIR}$ of NGC 7252 is about $10^{-4}$, and find that the position of a closed circle in Figure 3 presenting NGC 7252 is within the region of starburst galaxies. The luminosity ratio suggests that the soft X-ray emission is associated with the starburst activity. Note that the X-ray luminosity of NGC 7252 includes X-ray emissions from both soft ($L_{\rm X} = 2 \times 10^{40}$) and hard ($L_{\rm X} = 2.7 \times 10^{40}$) components, because Canizares et al. (1987) and David et al. (1992) did not distinguish soft and hard components when they estimated the X-ray luminosities.

In order to estimate the total amount of the hot gas and the total energy, we obtained the volume emission measure, $\varepsilon = \int n^2 dV$, of $(6.0^{+3.1}_{-2.0}) \times 10^{62}$ cm$^{-3}$ from the spectral fit of the X-ray spectrum with solar abundance ($A_{\rm RS} = 1.0$). This emission measure is the lowest compared to the sample of early-type galaxies by Matsumoto et al. (1997). We note that the sample by Matsumoto et al. (1997) have a mean abundance of 0.3, and that the volume emission measure of a thin thermal plasma is a strong function of the metal abundance. We obtained the emission measure $\varepsilon \sim 1.9 \times 10^{63}$ cm$^{-3}$ in $A_{\rm RS}$=0.3, and confirmed that the emission measure is still low among their sample. Read & Ponman (1998) found that the *ROSAT* PSPC source appeared centrally concentrated and compact, though a couple of tentative extended features were seen. The X-ray image by the *ROSAT* observation is consistent with the H$\alpha$ image reported by H94. In starburst galaxies, the X-ray emitting region is well correlated with the H$\alpha$ emitting region (for example, NGC 1808 (Dahlem, Hartner, & Junkes 1994)). Therefore, we assumed that the hot gas fills the inner region of $r$=2 kpc with a filling factor $f$, where $r$ is a radius of the hot plasma, because H94 pointed out that about 90 % of the H$\alpha$ line emission is in $r < 7''$ ($\sim$ 2.5 kpc at D = 63.2 Mpc). We obtain the mean electron number density to be $< n_e > \sim 2 \times 10^{-2} (r/2{\rm kpc})^{-3/2} f^{-1/2}$ cm$^{-3}$. The total mass and the total energy are estimated to be $2 \times 10^7 (r/2{\rm kpc})^{3/2} f^{1/2} M_{\odot}$ and $7 \times 10^{55} (r/2{\rm kpc})^{3/2} f^{1/2}$ ergs, respectively. They indicate that in starburst activity, NGC 7252 can stand comparison with nearby active starburst galaxies (e.g. M82 and NGC 253 (Fabbiano 1988) ).

Ciotti et al. (1991) classified the early-type galaxies into three evolutionary stages: wind, outflow, and inflow phases. Since it is thought that the intense star formation occurred within the last 1 Gyr (Schweitzer 1982; Hibbard & van Gorkom 1996) and numerical simulations have suggested that NGC 7252 evolves to an early-type galaxy, we consider that NGC 7252 belongs to the wind phase, in which the thermalized gas due to collisions between stellar winds and/or between winds and ambient gas is ejected from the galaxy. The low X-ray luminosity of NGC 7252 can be explained by the gas ejection from the galaxy. Based on the evolutional model by Ciotti et al. (1991), within a few Gyr, NGC 7252 will change to outflow phase, and then NGC 7252 will be surrounded by a large amount of hot gas.

### 3.2. Hard Component

Read & Ponman (1998) suggested the existence of hard X-ray sources in NGC 7252, since the fitted absorbing column appears larger than the Galactic value. The *ASCA*'s wide-band spectroscopy resolved the hard X-ray emission with the X-ray luminosity of about $5.6\times10^{40}$ ergs $\mathrm{s}^{-1}$ in the 2–10 keV band. The spectrum is characterized by the thermal model with $kT > 5.1$ keV or the power law model with photon index of 1.2.

The *ROSAT* HRI found two bright X-ray sources (Read & Ponman 1998). One is located within $7''$ from the center of NGC 7252, and its X-ray luminosity is estimated to be $3\times10^{40}$ ergs $\mathrm{s}^{-1}$ assuming the thermal emission with $kT = 3$ keV. The other is located in the eastern extension seen in the PSPC, and its X-ray luminosity is estimated to be $1.6\times10^{40}$ ergs $\mathrm{s}^{-1}$. *ASCA* cannot separate these two sources due to its poor spatial resolution. Since the X-ray luminosity observed by *ASCA* is nearly equal to the sum of the X-ray luminosities of the two sources, we consider that most of the hard X-ray emission come from these two sources. From the brightness of these two sources in the *ROSAT* HRI band, the significant fraction of the hard X-rays detected with *ASCA* is probably emitted from the former bright HRI source. Our result indicates the existence of a hard X-ray source with $L_{\mathrm{X}} \sim 10^{40}$ erg $\mathrm{s}^{-1}$ in the central region. The hard X-ray source may be a low-luminosity AGN or an intermediate-mass black hole ($10^{3-6}M_{\odot}$) (Colbert & Mushotzky 1999; Matsumoto et al. 2001; Kaaret et al. 2001).

## 4. Conclusion

We analyzed *ASCA* data for a merging galaxy NGC 7252. The X-ray spectrum is well described by a two-component model, soft plus hard components. The soft component is characterized by a thin thermal plasma model with $kT$ ~0.73keV. From the $L_{\mathrm{X}}/L_{\mathrm{FIR}}$, the thin thermal emission probably originates from the hot gas due to star formation. Assuming the size of hot gas of 2 kpc, the total amount of hot gas is estimated to be $2\times10^{7}f^{1/2}M_{\odot}$, which indicate that NGC 7252 still have large starburst activity comparable to those of nearby active starburst galaxies. Numerical simulations and observational results suggest that this galaxy is in the post-starburst phase. Therefore, we can explain the low X-ray luminosity of NGC 7252 due to ejection of hot gas from the galaxy. Based on the evolutional model by Ciotti et al. (1990), this galaxy will be surrounded by a large amount of hot gas within a few Gyr.

We resolve the hard X-ray emission characterized by the thermal emission with $kT > 5.1$ keV or a power law emission with $\Gamma$ ~1.2 using the *ASCA*'s wide-band spectroscopy. The *ROSAT* HRI found two bright point sources in this galaxy. These sources are possible objects for the hard X-ray emission. Our *ASCA* and the *ROSAT* results may suggest the existence of a low-luminosity AGN or an intermediate-mass black hole.

This galaxy is one of the best targets to investigate the evolution of merging galaxies and the connection between starburst galaxies and AGNs. The *XMM-Newton*, *Chandra*, and future missions will solve the current problems.

The authors thank all members of the *ASCA* team, and the staff in the *ASCA* GOF for providing *ASCA* data. We thank Ms. D. Gage and Ms. M. Awaki for their help. We thank the staff in the SkyView data surveys for providing the digitized sky survey image of NGC 7252. This research is supported by JSPS (HM), and is carried out as part of the "Ground Research Announcement for Space Utilization" promoted by the Japan Space Forum (HA).

## REFERENCES

Awaki, H., et al. 1994, PASJ, 46, L65

Bregman, J. N., Snider, B. A., Grego, L., & Cox, C.V. 1998, ApJ, 499, 670

Canizares, C. R., Fabbiano, G., & Trinchieri, G. 1987, ApJ, 312, 503

Ciotti, L., D'Ercole, A., Pellegrini, S., & Renzini, A. 1991, ApJ, 376, 380

Colbert, E. J. M., & Mushotzky, R. F. 1999, ApJ, 519, 89

Dahlem, M., Hartner, G. D., & Junkes, N. 1994, ApJ, 432, 598

David, L. P., Jones, C., & Forman, W. 1992, ApJ, 388, 82

Devereux, N. A, & Eales, S. A. 1989, ApJ, 340, 708

Dupraz, C., Casoli, F., Combes, F., & Kazes, I. 1990, A&A, 228, L5

Fabbiano, G. 1988, ApJ, 330, 672

Fabbiano, G., & Schweizer, F. 1995, ApJ, 447, 572

Forman, W., Jones, C., & Tucker, W. 1985, ApJ, 293, 102

Hibbard, J. E., Guhathakurta, P., van Gorkom, J. H., & Schweizer, F. 1994, AJ, 107, 67 (H94)

Hibbard, J. E., & van Gorkom, J. H. 1996, AJ, 111, 655

Hibbard, J. E., & Mihos, J. C. 1995, AJ, 110, 140

Kaaret, P., Prestwich, A. H., Zezas, A., Murray, S. S., Kim, D. -W., Kilgard, R. E., Schlegel, E. M., & Ward, M. J. 2001, MNRAS, 321, L29

Knapp, G. R., Guhathakurta, P., Kim, D. -W., & Jura, M. 1989, ApJS, 70, 329

Matsumoto, H., Koyama, K., Awaki, H., Tsuru, T., Loewenstein, M., & Matsushita, K. 1997, ApJ, 482, 133

Matsumoto, H., et al. 2001, ApJ, 547, L25

Matsushita, K., Ohashi, T., & Makishima, K. 2000, PASJ, 52, 685

Mihos, J. C., Dubinski, J., & Hernquist, L. 1998, ApJ, 494, 183

Read, A. M., & Ponman, T. J. 1998, MNRAS, 297, 143

Schweitzer, F. 1982, ApJ, 252, 455

Stanford, S. A., & Bushouse, H. A. 1991, ApJ, 371, 92

Stark, A.A., Gammie, C. F., Wilson, R. W., Bally, J., Linke, R. A., Heiles, C., & Hurwitz, M. 1992, ApJS, 79, 77

Tanaka, Y., Inoue, H., & Holt, S. S. 1994, PASJ, 46, L37

Whitemore, B. C., Miller, B. W., Schweizer, F., & Fall, S. M. 1997, AJ, 114, 1797

---

This 2-column preprint was prepared with the AAS LATEX macros v4.0.

# TABLE 1

# RESULT OF SPECTRAL FIT

| Model[a] | $N_{\mathrm{H}}^{b}$ ($\times 10^{22}$ cm$^{-2}$) | $kT_{\mathrm{RS}}$ (keV) | $A_{\mathrm{RS}}$ (solar) | $kT_{\mathrm{BR}}/\Gamma$ (keV) | $\chi^2_\nu$/d.o.f. |
|---|---|---|---|---|---|
| RS + BR | 0.02(fixed) | 0.73 (0.60-0.86) | 5.0 (0.05< ) | 4.5 < | 1.24/13 |
| RS + BR | 0.02(fixed) | 0.72 (0.61-0.85) | 1.0 (fixed) | 5.1 < | 1.16/14 |
| RS + PL | 0.02(fixed) | 0.73 (0.60-0.86) | 5.0 (0.03<) | 1.21(0.45-1.88) | 1.25/13 |
| RS + PL | 0.02(fixed) | 0.73 (0.60-0.86) | 1.0 (fixed) | 1.21(0.72-1.86) | 1.16/14 |

[a] the RS + BR indicates a Raymond-Smith optically thin plasma (RS) plus a bremsstrahlung (BR) model, and the RS + PL indicates a RS plus a power law (PL) model.

[b] the column density is fixed at the Galactic absorbing column density given in Stark et al. (1992).

NOTE.—errors are 90 % confidence level.



Fig. 1.— *ASCA* SIS contour map overlaid the digitized sky survey image of NGC 7252. The map has been smoothed with $\sigma$=30″. The contours are at 2, 3, 4, and 5 $\sigma$ fluctuation levels of the SIS background.



Fig. 2.— *ASCA* SIS spectrum of NGC 7252 fitted with a two-component model, consisting of a soft RS plasma component (dashed line) and a hard bremsstrahlung component (dot-dashed line).



Fig. 3.— Correlation plot between far-infrared and the 0.5–4.5 keV band luminosities. Open circles and one closed circle show the data for early-type galaxies and for NGC 7252, respectively. The region between two solid lines indicates the region of spiral galaxies found by David et al. (1991). A couple of early-type galaxies locate in the region of spiral galaxies, but they are also classified as Seyfert/LINER.